WEST Search History

DATE: Friday, February 28, 2003

Set Name Query side by side			Hit Count Set Name result set		
DB=U	SPT,PGPB,JPAB,EPAB,DWPI; THES=ASSIGNEE; PLUR=YES;				
OP=ADJ					
L13	L10 same (therap\$ or vivo or administ\$)	56	L13		
L12	L10 same oralS	3	L12		
1.11	L10 same (11 or 12 or 13 or 14 or 15 or 16 or 17 or 18)	()	L11		
L10	L9 same (igg or immunoglobulin or antibody)same (eggS or milk or plasma or blood)	180	L10		
L9	chronic fatigue syndrome or cfs	87960	L9		
DB = U	SPT: THES=ASSIGNEE; PLUR=YES; OP=ADJ				
L8	QUIGLEY-JAMES-S.in.	10	L8		
L7	ARTHINGTON-JOHN-S.in.	()	L7		
L6	POLO POZO-FRANCISCO-S	()	L6		
L5	RUSSELL-LOUIS-\$.in.	1	L5		
L4	BORG-BARTON-S.in.	()	1.4		
L3	WEAVER-ERIC-S.in.	3	L3		
L.2	STROHBEHN-RONALD-S.in.	()	L2		
L1	CAMPBELL-JOY-S.in.	1	L1		

END OF SEARCH HISTORY

d his

(FILE 'HOME' ENTERED AT 09:45:47 ON 28 FEB 2003)

F:LE 'CAPLUS, 1MOBILITY, AGRICOLA, AQUASCI, BIOTECHNO, COMPENDEX, COMPUAB, CONF, CONFSCI, ELCOM, EVENTLINE, HEALSAFE, IMSDRUGGONF, LIFESCI, OCEAN, MEDICONF, PASCAL, PAPERCHEM2, POLLUAB, SOLIDSTATE, ADISCTI, ADISINSIGHT, ADISNEWS, ANABSTR, BIOBUSINESS, ...' ENTERED AT 09:46:20 ON 28 FEB 2003

	28	FEB	200	3	
				Ε	CAMPBELL JOY?/AU
L1			20	S	E1 OR E2
				Ε	STROHBEHN RONALD?/AU
L2			11	S	E1 OR E2
				Ε	WEAVER ERIC?/AU
L3			15	S	E2 OR E1
				Ε	BORG BARTON?/AU
L4			2	S	E2
				Е	RUSSELL LOUIS?/AU
L5			13	S	E1 OR E2
				Ε	POLO POZO FRANCISCO?/AU
L6			6	S	E1 OR E2
				Ε	ARTHINGTON JOHN?/AU
L7			5	S	El OR E4
				Ε	QUIGLEY JAMES?/AU
L8			18	S	E1 OR E2
L9		25	5304	S	CHRONIC FATIGUE SYNDROME OR CFS
L10			457	S	L9 (S) (IGG OR IMMUNOGLOBULIN? OR ANTIBOD?) (S) (EGG OR MIL
L11			87	S	L10 (S) (VIVO OR ADMINISTER? OR THERAP?)
L12			51	Dt	JP REM L11 (36 DUPLICATES REMOVED)

12 ANSWER 45 OF 51 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.DUPLICATE

ACCESSION NUMBER: 1991:53089 BIOSIS

BA91:31370 DOCUMENT NUMBER:

A CONTROLLED TRIAL OF INTRAVENOUS IMMUNOGLOBULIN G IN TITLE:

CHRONIC FATIGUE SYNDROME.

PETERSON P K; SHEPARD J; MACRES M; SCHENCK C; CROSSON J; AUTHOR(S):

RECHTMAN D; LURIE N

DEP. OF MED., HENNEPIN COUNTY MED. CENT., 701 PARK AVE., CORPORATE SOURCE:

MINNEAPOLIS, MINN. 55415.

AM J MED, (1990) 89 (5), 554-560. SOURCE:

CODEN: AJMEAZ. ISSN: 0002-9343.

FILE SEGMENT: BA; OLD LANGUAGE: English

PURPOSE: Currently, there is no established therapy for

chronic fatigue syndrome (CFS), a

recently defined illness that has been associated with a variety of immunologic abnormalities. Based on the hypothesis that a chronic viral infection or an immunoregulatory defect is involved in the pathogenesis of

CFS, the therapeutic benefit of intravenous immunoglobulin G (IV IgG) was evaluated in a group of patients with CFS. Additionally, serum immunoglobulin concentrations and peripheral **blood** lymphocyte subset numbers were measured at the outset of the study, and the effect of IV IgG therapy on IgG subclass levels was determined. PATIENTS AND METHODS: Thirty patients with CFS were enrolled in a double-blind, placebo-controlled trial of IV IgG. The treatment of regimen consisted of IV IgG (1 g/kg) or intravenous placebo (1% albumin solution) administered every 30 days for 6 months. Participants completed a self-assessment form prior to each of the six treatments, which was used to measure severity of symptoms, functional status, and health perceptions. Patients were also asked to report adverse experiences defined as worsening of symptoms occurring within 48 hours of each treatment. RESULTS: Twenty-eight patients completed the trial. At baseline, all 28 patients complained of moderate to severe fatigue, and measures of social functioning and health perceptions showed marked impairment. Low levels of IgG1 were found in 12 (42.9%), and 18(64.3%) had low levels of IgG3. At the end of the study, no significant therapeutic benefit could be detected in terms of symptom amelioration or improvement in functional status, despite restoration of

IgG1 levels to a normal range. Major adverse experiences were observed in 20% of both the IV IqG and placebo groups. CONCLUSION: The results of this study indicate that IV IgG is unlikely to be of clinical benefit in CFS. In addition to the ongoing need for placebo controlled trials of candidate therapies for CFS

, an expanded research effort is needed to define the etiology